

## Switching the topological charge of exciton polariton vortices

Berger, Bernd<sup>1,\*</sup>, Ma, Xuekai<sup>2</sup>, Schneider, Christian<sup>3</sup>, Höfling, Sven<sup>3,4</sup>, Schumacher, Stefan<sup>2,5</sup>, Assmann, Marc<sup>1</sup>

<sup>1</sup>Experimentelle Physik 2, Technische Universität Dortmund, 44227 Dortmund, Germany

<sup>2</sup>Department of Physics and Center for Optoelectronics and Photonics Paderborn (CeOPP), Universität Paderborn, Warburger Strasse 100, 33098 Paderborn, Germany

<sup>3</sup>Technische Physik, Universität Würzburg, Am Hubland, 97074, Würzburg, Germany


<sup>4</sup>SUPA, School of Physics and Astronomy, University of St. Andrews, St. Andrews KY16 9SS, United Kingdom

<sup>5</sup>College of Optical Sciences, University of Arizona, Tucson, AZ 85721, USA

\*e-mail: bernd.berger@tu-dortmund.de

Vortices are elementary excitations arising in exciton polariton condensates [1] and consist of a  $2\pi \cdot m$  radial phase shift of the polariton wavefunction, where  $m$  is the topological charge of the vortex. This topological charge translates into the orbital angular momentum (OAM) state of the emitted light field and can be detected by application of a dedicated OAM spectroscopy technique [2]. This technique gives us the opportunity to study the temporal dynamics of vortices in exciton polariton condensates without using complex interferometric techniques that also always require a phase reference. Here, we experimentally demonstrate the switching of the topological charge of a vortex due to a pulsed perturbation [3].

First, we create a stable localized  $m=+1$  vortex inside an annular trap created all-optically by a CW laser. Then an additional non-resonant pulsed laser is switched on in addition to create a potential barrier, which interacts with the polariton vortex and may reverse its direction of rotation. Depending on the pumping power of the pulsed laser a switching to the  $m=-1$  state is observed. This effect is shown in Fig. 1.



**Fig. 1.** Left: Vortex switching in dependence of the excitation power of the pulsed perturbation. Right: Experimental geometry.

### References

- [1] K. G. Lagoudakis, M. Wouters, M. Richard, A. Baas, I. Carusotto, R. André, Le Si Dang and B. Beveaud-Plédran, Nat. Phys., 2008, **4**, 706-710.
- [2] B. Berger, M. Kahlert, D. Schmidt and M. Assmann, Opt. Express, 2018, **26**, 32248-32258.
- [3] X. Ma, B. Berger, M. Assmann, R. Driben, T. Meier, C. Schneider, S. Höfling and S. Schumacher, arXiv:1907.03171, submitted 6 July, 2019.