

BEC-BCS crossover in an equilibrium exciton-polariton condensate

Hu, Hui¹ and Liu, Xia-Ji^{1*}

¹*Centre for Quantum and Optical Science, Swinburne University of Technology, Melbourne, Victoria 3122, Australia*

**e-mail: xiajiliu@swin.edu.au*

Microcavity exciton-polariton systems in two-dimensions are long anticipated to exhibit a crossover from Bose-Einstein condensate (BEC) to Bardeen-Cooper-Schrieffer (BCS) superfluid, when the carrier density is tuned to reach the Mott transition density. Yet, theoretical understanding of such a BEC-BCS crossover largely relies on the mean-field framework [1, 2] and the nature of carriers at the crossover remains unclear to some extent. Here, inspired by the recent demonstration of a BCS polariton laser [3] and based on a simplified description with contact interactions to model the attraction between electrons and holes, we investigate quantum fluctuations of an exciton-polariton condensate and determine the number of electron-hole pairs, condensed photons and polaritons at the crossover beyond mean-field. We find that the exciton-polariton condensate remains in the strong-coupling regime, when the carrier density is much larger than the Mott density.

References

- [1] K. Kamide and T. Ogawa, Phys. Rev. Lett. 105, 056401 (2010).
- [2] T. Byrnes, T. Horikiri, N. Ishida, and Y. Yamamoto, Phys. Rev. Lett. 105, 186402 (2010).
- [3] J. Hu, Z. Wang, S. Kim, H. Deng, S. Brodbeck, C. Schneider, S. Höfling, N. H. Kwong, and R. Binder, Signatures of a Bardeen-Cooper-Schrieffer Polariton Laser, arXiv:1902.00142v1.